



TITLE:

行列環のテンソル積空間上の部分 転置写像について (線形作用素に関 連する不等式とその応用)

AUTHOR(S):

SANO, Takashi

CITATION:

SANO, Takashi. 行列環のテンソル積空間上の部分転置写像について (線形作用素に関連する不等式とその応用). 数理解析研究所講究録 2008, 1596: 140-142

ISSUE DATE:

2008-04

URL:

<http://hdl.handle.net/2433/81698>

RIGHT:

行列環のテンソル積空間上の部分転置写像について

山形大学・理学部数理科学科 佐野隆志 (Takashi SANO)
Department of Mathematical Sciences
Faculty of Science, Yamagata University

This is a survey of [1], which is a joint work with Tsuyoshi Ando. Here are some of the results in [1]. For more details, we refer the reader to [1].

1. 部分転置写像とその関連

Theorem 1.1. *Let $A_k \in M_m$, $B_k \in M_n$. Then for every unitarily invariant norm $||| \cdot |||$*

$$\sup_{X \in M_{m,n}} \frac{||| \sum_k A_k X B_k^T |||}{||| X |||} \leq \sqrt{\min(m, n)} \left\| \sum_k A_k \otimes B_k \right\|_{\infty}.$$

Theorem 1.2. *Let $A_k \in M_m$, $B_k \in M_n$. Then for every unitarily invariant norm $||| \cdot |||$*

$$\sup_{X \in M_{m,n}} \frac{||| \sum_k A_k X B_k |||}{||| X |||} \leq \min(m, n) \left\| \sum_k A_k \otimes B_k \right\|_{\infty}.$$

The partial transpose map Θ on $M_m \otimes M_n$ is defined as

$$\Theta\left(\sum_k A_k \otimes B_k\right) = \sum_k A_k \otimes B_k^T.$$

Theorem 1.3. *For every unitarily invariant norm $||| \cdot |||$*

$$||| \Theta(X) ||| \leq \min(m, n) ||| X |||.$$

For the Hilbert-Schmidt norm $\| \cdot \|_2$

$$\| \Theta(X) \|_2 = \| X \|_2.$$

Notice that this inequality for the spectral norm was originally established by Tomiyama

2. 部分転置写像の制限

Let

$$K_{m,n} := \{A \otimes I_n + I_m \otimes B : A \in M_m, B \in M_n\}.$$

We observe the restriction Θ_0 of the map Θ to the subspace $K_{m,n}$:

$$\Theta_0(X) = A \otimes I_n + I_m \otimes B^T \quad \text{for } X = A \otimes I_n + I_m \otimes B.$$

Theorem 2.1. *The restriction Θ_0 is positive:*

$$X \geq 0 \implies \Theta_0(X) \geq 0.$$

Theorem 2.2. *Let $X = A \otimes I_n + I_m \otimes B$.*

(i) *If $n = 2$ or B is normal, then X and $\Theta_0(X)$ are unitarily similar in $M_m \otimes M_n$.*

(ii) *If $m = 2$ or A is normal, then X and $\Theta_0(X)^T$ are unitarily similar in $M_m \otimes M_n$.*

In any case, $|X|$ and $|\Theta_0(X)|$ are unitarily similar in $M_m \otimes M_n$; hence, $|||\Theta_0(X)||| = |||X|||$ for every unitarily invariant norm $|||\cdot|||$.

Theorem 2.3. *If $X \in K_{m,n}$ is normal, then X and $\Theta_0(X)$ are unitarily similar and $|||\Theta_0(X)||| = |||X|||$ for every unitarily invariant norm $|||\cdot|||$.*

Theorem 2.4. *For every unitarily invariant norm $|||\cdot|||$*

$$|||\Theta_0(X)||| \leq 2 |||X||| \quad (X \in K_{m,n}).$$

We recall the definition of the *numerical range* $W(X)$ of a square matrix $X \in M_n$:

$$W(X) := \{\langle Xa, a \rangle : a \in \mathbb{C}^n, \|a\| = 1\}$$

The *numerical radius* $w(X)$ is defined as

$$w(X) = \sup\{|\xi| : \xi \in W(X)\} = \sup\{|\langle Xa, a \rangle| : a \in \mathbb{C}^n, \|a\| = 1\}$$

Theorem 2.5. *For $X \in K_{m,n}$*

$$w(\Theta_0(X)) = w(X) \quad \text{and more precisely, } W(\Theta_0(X)) = W(X).$$

Theorem 2.6. *Let $A \in M_m, B \in M_n$. Then for every unitarily invariant norm $|||\cdot|||$*

$$\sup_{X \in K_{m,n}} \frac{|||AX + XB^T|||}{|||X|||} \leq \sqrt{2} \|A \otimes I_n + I_m \otimes B\|_\infty,$$

and

$$\sup_{X \in K_{m,n}} \frac{|||AX + XB|||}{|||X|||} \leq \sqrt{2} \|A \otimes I_n + I_m \otimes B\|_\infty.$$

Theorem 2.7. For $A \in M_m$ and $B \in M_n$

$$\|A \otimes I_m + I_n \otimes B^T\|_\infty \leq \sqrt{2} \|A \otimes I_n + I_m \otimes B\|_\infty.$$

Recall the definition of the *Schatten p -norm* $\|\cdot\|_p$ for $1 \leq p < \infty$:

$$\|X\|_p := \left\{ \sum_i s_i(X)^p \right\}^{1/p}.$$

Theorem 2.8. For $p = 2k$ ($k = 1, 2, 3, 4, 5$)

$$\|\Theta_0(\mathbf{X})\|_p = \|\mathbf{X}\|_p \quad (\mathbf{X} \in K_{m,n}).$$

REFERENCES

- [1] T. Ando and T. Sano, *Norm estimates of the partial transpose map on the tensor products of matrices*, Positivity, 12 (2008), 9-24.